



FLUID ORGANIC NITROGEN FERTILISER N 3%
BORON 6% COMPLEXED WITH ETHANOLAMINE



STRENGTHS

- SPECIFIC FOR PREVENTING AND RESOLVING BORON DEFICIENCIES
- FACILITATES THE TRANSLOCATION OF BORON WITHIN PLANTS
- IMPROVES FRUIT SETTING IN OLIVES, GRAPEVINES, KIWIS, FRUIT TREES AND VEGETABLES
- IMPROVES PHOTOSYNTHETIC, GERMINATION AND ROOTING PROCESSES
- STIMULATES THE ELONGATION OF NEW SHOOTS IN THE OLIVE TREE BY AVOIDING PRODUCTION ALTERNATION
- PROMOTES GREATER VEGETATIVE AND PRODUCTIVE DEVELOPMENT



Allowed
in organic
farming



PACKAGE

Bottles 1 L (20x1)
Jerrycans 5 L (4x5)



CHEMICAL-PHYSICAL PROPERTIES

Formulation: **liquid**
Density: **1,430**
pH (sol.1%): **8.5 ± 1**
Conductivity (1‰):
mS/cm: **0.17**



FEATURES

BOROLIVO Plus is a product based on organic Nitrogen (N) and Boron complexed with ethanolamine. This unique product is rich in amino acids and sugars which, in addition to their humectant function, once inside the leaves, facilitate the penetration of boron into the plants, making it particularly effective on all crops. Encourages the pollinating behaviour of hymenopterous insects (bees and bumblebees).

BOROLIVO Plus is therefore the specific product to deal with the following issues:

- POOR FRUIT SETTING IN OLIVE TREES, VINES, KIWI, FRUIT AND VEGETABLES
- MILLERANDAGE IN VINE
- SUBEROSITY OF FRUITS (PEAR AND APPLE TREE)
- HOLLOW HEART OF THE SUGAR BEET
- IMPROVEMENT OF PHOTOSYNTHETIC, GERMINATIVE AND ROOTING PROCESSES



DOSES AND METHODS OF USE

CROPS

FOLIAR APPLICATION

Grapevine (reduction of millerandage)
Kiwi

100-150 ml/hl pre-flowering and post setting

Olive - Citrus - Date Palm

200-250 ml/hl pre-flowering,
flowering and post setting

Fruit crops:
(Pear, Apple, Peach, Plum, Cherry, Apricot, etc.)

100-150 ml/hl pre-flowering and
post setting

Horticultural crops in greenhouses and open fields:
(Tomato, Pepper, Aubergine, Courgette, Melon,
Cucumber, Strawberry, Bean, Asparagus etc.)

100-150 ml/hl for recognised needs only

**Sugar beet, Leafy and cut vegetables,
Aromatics Cauliflower,
Carrot, Sunflower, Potato, Turnip,**

150-200 ml/hl at the 4-6 leaf stage

Legumes, Floricultural, Forestry

150-200 ml/hl preventively or as needed

COMPOSITION

Organic nitrogen (N)	3,0%
Water-soluble organic nitrogen (N)	3,0%
Organic carbon (C)	10,0%
Water-soluble boron (B)	6,0%



Use only in cases of recognised need.

Do not exceed the appropriate doses.

Combination with Foliar PH (acidifier - humectant

for solutions) 100 g/hl is recommended.

Do not mix BOROLIVO with white or activated oils.

DO NOT MIX WITH ALUMINIUM-BASED PRODUCTS (Al)

