



FERTIRAME 20

CE FERTILISER – MICRO-NUTRIENT MIXTURE
COPPER (Cu) 20% (OXYCHLORIDE) and BORON (B) 0.5%



STRENGTHS

- THE PRESENCE OF **BORON** HELPS PREVENT VARIOUS PHYSIOPATHOLOGIES SUCH AS MILLERANDAGE IN VINES AND POOR FRUIT SET IN OLIVE TREES
- ROCK POWDER PROTECTS PLANT ORGANS AND FRUIT FROM TEMPERATURE CHANGES AND BURNS
- ITS CONSTITUENTS AS A WHOLE HAVE A REPELLENT ACTION
- STRENGTHENS THE PLANTS' NATURAL DEFENCES AGAINST BIOTIC ADVERSITIES
- COPPER AND ZEOLITES "CAMOUFLAGE" THE DRUPE AND DISCOURAGE OLIVE FLY OVIPOSITION



Allowed in
organic farming



PACKAGE

Bags
1 - 3 - 10 kg



FEATURES

FERTIRAME 20 is a formulation containing **Copper (Oxychloride)** and **Boron (B)** with **Rock Powders (Zeolite and Kaolin)**. These constituents prevent and treat physiopathic symptoms caused by deficiencies of these micro-nutrients in many crops, strengthen defences against biotic adversities and, thanks to the presence of rock powder, create a coating that acts as a sunscreen, protecting plant organs and fruit from temperature fluctuations and sunburn caused by **UVA** and **UVB** rays; they are also known to have a natural repellent effect.

The presence of boron helps prevent the following pathologies:

- Desiccation of the rachis
- Grapevine millerandage
- Poor fruit setting in olives, fruit trees and vegetables
- Hollow heart of chard
- Shortening of internodes

DOSES AND METHODS OF USE

CROPS	APPLICAZIONE FOGLIARE
Viticulture	250-300 g/hl - (Normal Volume) - 2,5-3,0 kg/ha (Low Volume)
Olive, hazelnut, walnut, chestnut	250-350 g/hl - (Normal Volume) - 2,5-3,5 kg/ha (Low Volume)
Actinidia, apple, pear, peach and other stone fruits	Autumn-winter applications: 0,5-0,6 kg/hl (Normal Volume) 5-6 kg/ha (Low Volume)
Citrus fruits	250-350 g/hl - (Normal Volume) - 2,5-3,5 kg/ha (Low Volume)
Sugar beet, Carrot	250-350 g/hl - (Normal Volume) - 2,0-3,5 kg/ha (Low Volume)
Potato, tomato, pepper, courgette, garlic and onion	200-300 g/hl - (Normal Volume) - 2,0-3,0 kg/ha (Low Volume)
Artichoke, cabbage, chicory, pea, spinach, melon, watermelon, cucumber, fennel, radish	250-300 g/hl - (Normal Volume) - 2,5-3,0 kg/ha (Low Volume)
Strawberry	Sprouting: 200-300 gr/hl (V.N.) - 2,0-3,0 kg/ha (Low Volume)
Leafy-cut vegetables, Fresh-cut leaf, baby leaf, Lettuce, Endives, Aromatic crops	150-200 g/hl - (Normal Volume) - 1,5-2,0 kg/ha (Low Volume)
Rice, Wheat, Grains and Leguminous plants	200-300 g/hl - (Normal Volume) - 2,0-3,0 kg/ha (Low Volume)
Plants in nurseries	200 g/hl - (Normal Volume) - 2,0 kg/ha (Low Volume)

N.B. Do not mix with products with alkaline reaction. Do not mix with organic products and in particular with protein hydrolysates. Before mixing with crop protection agents or other fertilisers, carry out preliminary compatibility tests. Do not use products during flowering on pome and stone fruit (peach, plum, apple, pear...) In greenhouses: do not exceed 2 kg/ha

COMPOSITION

Water-soluble boron (B)	0,5%
Copper (Cu) total	20,0%

Copper (Cu) Oxchloride 20%



CHEMICAL-PHYSICAL PROPERTIES

Formulation: **Wettable powder**
 Colour: Blue - pH (sol. 1%): **7.7 ±1**
 Conductivity (1‰) mS/cm 18° C: **0.80**
 Solubility: **30 g/100 ml**

To be used only in cases of recognised need.
Do not exceed the appropriate doses



Before use, carefully read the hazard (H) statements on page 172.

